

Changing Variables in Multiple Integrals

1. Changing variables.

Double integrals in x, y coordinates which are taken over circular regions, or have integrands involving the combination $x^2 + y^2$, are often better done in polar coordinates:

$$(1) \quad \iint_R f(x, y) dA = \iint_R g(r, \theta) r dr d\theta .$$

This involves introducing the new variables r and θ , together with the equations relating them to x, y in both the forward and backward directions:

$$(2) \quad r = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1}(y/x); \quad x = r \cos \theta, \quad y = r \sin \theta .$$

Changing the integral to polar coordinates then requires three steps:

- A. Changing the integrand $f(x, y)$ to $g(r, \theta)$, by using (2);
- B. Supplying the area element in the r, θ system: $dA = r dr d\theta$;
- C. Using the region R to determine the limits of integration in the r, θ system.

In the same way, double integrals involving other types of regions or integrands can sometimes be simplified by changing the coordinate system from x, y to one better adapted to the region or integrand. Let's call the new coordinates u and v ; then there will be equations introducing the new coordinates, going in both directions:

$$(3) \quad u = u(x, y), \quad v = v(x, y); \quad x = x(u, v), \quad y = y(u, v)$$

(often one will only get or use the equations in one of these directions). To change the integral to u, v -coordinates, we then have to carry out the three steps **A, B, C** above. A first step is to picture the new coordinate system; for this we use the same idea as for polar coordinates, namely, we consider the grid formed by the level curves of the new coordinate functions:


$$(4) \quad u(x, y) = u_0, \quad v(x, y) = v_0 .$$

Once we have this, algebraic and geometric intuition will usually handle steps **A** and **C**, but for **B** we will need a formula: it uses a determinant called the **Jacobian**, whose notation and definition are

$$(5) \quad \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} x_u & x_v \\ y_u & y_v \end{vmatrix} .$$

Using it, the formula for the area element in the u, v -system is

$$(6) \quad dA = \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv ,$$



so the change of variable formula is

$$(7) \quad \iint_R f(x, y) \, dx \, dy = \iint_R g(u, v) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv ,$$

where $g(u, v)$ is obtained from $f(x, y)$ by substitution, using the equations (3).

We will derive the formula (5) for the new area element in the next section; for now let's check that it works for polar coordinates.

Example 1. Verify (1) using the general formulas (5) and (6).


Solution. Using (2), we calculate:

$$\frac{\partial(x, y)}{\partial(r, \theta)} = \begin{vmatrix} x_r & x_\theta \\ y_r & y_\theta \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r(\cos^2 \theta + \sin^2 \theta) = r ,$$

so that $dA = r \, dr \, d\theta$, according to (5) and (6); note that we can omit the absolute value, since by convention, in integration problems we always assume $r \geq 0$, as is implied already by the equations (2).

We now work an example illustrating why the general formula is needed and how it is used; it illustrates step **C** also — putting in the new limits of integration.

Example 2. Evaluate $\iint_R \left(\frac{x-y}{x+y+2} \right)^2 \, dx \, dy$ over the region R pictured.



Solution. This would be a painful integral to work out in rectangular coordinates. But the region is bounded by the lines

$$(8) \quad x + y = \pm 1, \quad x - y = \pm 1$$

and the integrand also contains the combinations $x - y$ and $x + y$. These powerfully suggest that the integral will be simplified by the change of variable (we give it also in the inverse direction, by solving the first pair of equations for x and y):

$$(9) \quad u = x + y, \quad v = x - y; \quad x = \frac{u+v}{2}, \quad y = \frac{u-v}{2} .$$

We will also need the new area element; using (5) and (9) above. we get

$$(10) \quad \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{vmatrix} = -\frac{1}{2} ;$$

note that it is the second pair of equations in (9) that were used, not the ones introducing u and v . Thus the new area element is (this time we do need the absolute value sign in (6))

$$(11) \quad dA = \frac{1}{2} \, du \, dv .$$

We now combine steps **A** and **B** to get the new double integral; substituting into the integrand by using the first pair of equations in (9), we get

$$(12) \quad \iint_R \left(\frac{x-y}{x+y+2} \right)^2 \, dx \, dy = \iint_R \left(\frac{v}{u+2} \right)^2 \frac{1}{2} \, du \, dv .$$

In uv -coordinates, the boundaries (8) of the region are simply $u = \pm 1$, $v = \pm 1$, so the integral (12) becomes

$$\iint_R \left(\frac{v}{u+2} \right)^2 \frac{1}{2} du dv = \int_{-1}^1 \int_{-1}^1 \left(\frac{v}{u+2} \right)^2 \frac{1}{2} du dv$$

We have

$$\text{inner integral} = \left. -\frac{v^2}{2(u+2)} \right]_{u=-1}^{u=1} = \frac{v^2}{3} ; \quad \text{outer integral} = \left. \frac{v^3}{9} \right]_{-1}^1 = \frac{2}{9} .$$

MIT OpenCourseWare
<http://ocw.mit.edu>

18.02SC Multivariable Calculus
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.