


# LECTURE 22

- Predict whether an aqueous solution of each of the following salts has a pH equal to, greater than or less than 7.  
(a)  $\text{KC}_6\text{H}_5\text{CO}_2$                       (c)  $\text{NH}_4\text{Br}$                                       (e)  $\text{Li}_2\text{CO}_3$   
(b)  $\text{NaNO}_2$                                       (d)  $\text{MgCl}_2$
- Phosphate buffers are very useful in biochemical experiments. Your UROP supervisor asks you to make up a phosphate buffer to carry out kinetic assays on an enzyme using the conjugate acid/base pair of  $\text{HPO}_4^{2-}$  ( $K_a = 2.1 \times 10^{-13}$ ) and  $\text{PO}_4^{3-}$ , both available from the stock room in the form of potassium salts.  
(a) What must be the ratio of the molarities of  $\text{PO}_4^{3-}$  and  $\text{HPO}_4^{2-}$  ions in a buffer solution having a pH of 12.0 (report ratio to one significant figure)?  
(b) What mass of  $\text{K}_3\text{PO}_4$  must be added to 2.00 L of 0.100 M  $\text{K}_2\text{HPO}_4(\text{aq})$  to prepare a buffer solution with a pH of 12.0 (report mass to one significant figure)?  
(c) State the range of pH values for which this phosphate buffer will be an effective in maintaining a constant pH.
- A different phosphate buffer is now put to test to see if it will maintain the pH of an enzyme solution if a strong base is added. This buffer solution was prepared to a final volume of 100.0 mL with concentrations of the salts of the conjugate acid/base pairs as following: 0.150 M  $\text{Na}_2\text{HPO}_4(\text{aq})$  and 0.100 M  $\text{KH}_2\text{PO}_4(\text{aq})$ . What are the pH and the pH change resulting from the addition of 80.0 mL of 0.0100 M  $\text{NaOH}(\text{aq})$  to the buffer solution? The  $\text{pK}_a$  of  $\text{H}_2\text{PO}_4^-$  is 7.21.
- A pharmaceutical molecule with antifungal properties is only active when deprotonated and negatively charged ( $\text{A}^-$ ). The protonated state ( $\text{HA}$ ) is inactive. If the  $\text{pK}_a$  of this drug is 9.0,  
(a) calculate the ratio of protonated to deprotonated compound at physiological pH (7.4).  
(b) Without doing a calculation, would more or less of the drug be active at  $\text{pH}=7.4$  if the  $\text{pK}_a$  of the drug was 8.0
- If 50.0 mL of a 0.200 M solution of the weak base *N*-ethylmorpholine ( $\text{C}_6\text{H}_{13}\text{NO}$ ) is mixed with 8.00 mL of 1.00 M  $\text{HCl}$  and then diluted to a final volume of 100.0 mL with water, the result is a buffer with a pH of 7.00. Compute the  $K_b$  of *N*-ethylmorpholine.
- Absorption of aspirin (acetylsalicylic acid,  $\text{C}_9\text{H}_8\text{O}_4$ ) into the bloodstream occurs only when the molecule is in its conjugate base form.  
(a) If a patient takes one tablet of aspirin (325 mg of aspirin), how many milligrams of aspirin are available for immediate absorption in the stomach? The pH of the stomach is 1.6, and the  $\text{pK}_a$  of aspirin is 3.5.  
(b) Would you expect more or less aspirin to be absorbed in the small intestine ( $\text{pH} \approx 7.5$ ) compared to the stomach? Briefly explain your answer (no calculation is required).



MIT OpenCourseWare  
<https://ocw.mit.edu>

5.111 Principles of Chemical Science  
Fall 2014

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.