


**Problem Set 6: Solutions**

1. Let us draw the region where  $f_{X,Y}(x,y)$  is nonzero:



The joint PDF has to integrate to 1. From  $\int_{x=1}^{x=2} \int_{y=0}^{y=x} ax \, dy \, dx = \frac{7}{3}a = 1$ , we get  $a = \frac{3}{7}$ .

$$(b) \quad f_Y(y) = \int f_{X,Y}(x,y) \, dy = \begin{cases} \int_1^2 \frac{3}{7}x \, dx, & \text{if } 0 \leq y \leq 1, \\ \int_y^2 \frac{3}{7}x \, dx, & \text{if } 1 < y \leq 2, \\ 0, & \text{otherwise} \end{cases} = \begin{cases} \frac{9}{14}, & \text{if } 0 \leq y \leq 1, \\ \frac{3}{14}(4 - y^2), & \text{if } 1 < y \leq 2, \\ 0, & \text{otherwise.} \end{cases}$$

(c)

$$f_{X|Y}(x \mid \frac{3}{2}) = \frac{f_{X,Y}(x, \frac{3}{2})}{f_Y(\frac{3}{2})} = \frac{8}{7}x, \quad \text{for } \frac{3}{2} \leq x \leq 2 \text{ and } 0 \text{ otherwise.}$$

Then,

$$\mathbf{E} \left[ \frac{1}{X} \mid Y = \frac{3}{2} \right] = \int_{3/2}^2 \frac{1}{x} \frac{8}{7}x \, dx = \frac{4}{7}.$$

(d) We use the technique of first finding the CDF and differentiating it to get the PDF.

$$\begin{aligned} F_Z(z) &= \mathbf{P}(Z \leq z) \\ &= \mathbf{P}(Y - X \leq z) \\ &= \begin{cases} 0, & \text{if } z < -2, \\ \int_{x=-z}^{x=2} \int_{y=0}^{y=x+z} \frac{3}{7}x \, dy \, dx = \frac{8}{7} + \frac{6}{7}z - \frac{1}{14}z^3, & \text{if } -2 \leq z \leq -1, \\ \int_{x=1}^{x=2} \int_{y=0}^{y=x+z} \frac{3}{7}x \, dy \, dx = 1 + \frac{9}{14}z, & \text{if } -1 < z \leq 0, \\ 1, & \text{if } 0 < z. \end{cases} \end{aligned}$$

$$f_Z(z) = \frac{d}{dz} F_Z(z) = \begin{cases} \frac{6}{7} - \frac{3}{14}z^2, & \text{if } -2 \leq z \leq -1, \\ \frac{9}{14}, & \text{if } -1 < z \leq 0, \\ 0, & \text{otherwise.} \end{cases}$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
 Department of Electrical Engineering & Computer Science  
**6.041/6.431: Probabilistic Systems Analysis**  
 (Fall 2010)

---

2. The PDF of  $Z$ ,  $f_Z(z)$ , can be readily computed using the convolution integral:

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t)f_Y(z-t) dt.$$

For  $z \in [-1, 0]$ ,

$$f_Z(z) = \int_{-1}^z \frac{1}{3} \cdot \frac{3}{4}(1-t^2) dt = \frac{1}{4} \left( z - \frac{z^3}{3} + \frac{2}{3} \right).$$

For  $z \in [0, 1]$ ,

$$f_Z(z) = \int_{z-1}^z \frac{1}{3} \cdot \frac{3}{4}(1-t^2) dt = \frac{1}{4} \left( 1 - \frac{z^3}{3} + \frac{(z-1)^3}{3} \right).$$

For  $z \in [1, 2]$ ,

$$f_Z(z) = \int_{z-1}^1 \frac{1}{3} \cdot \frac{3}{4}(1-t^2) dt + \int_{-1}^{z-2} \frac{2}{3} \cdot \frac{3}{4}(1-t^2) dt = \frac{1}{4} \left( z + \frac{(z-1)^3}{3} - \frac{2(z-2)^3}{3} - 1 \right).$$


For  $z \in [2, 3]$ ,

$$f_Z(z) = \int_{z-3}^{z-2} \frac{2}{3} \cdot \frac{3}{4}(1-t^2) dt = \frac{1}{6} (3 + (z-3)^3 - (z-2)^3).$$

For  $z \in [3, 4]$ ,

$$f_Z(z) = \int_{z-3}^1 \frac{2}{3} \cdot \frac{3}{4}(1-t^2) dt = \frac{1}{6} (11 - 3z + (z-3)^3).$$

A sketch of  $f_Z(z)$  is provided below.



3. (a)  $X_1$  and  $X_2$  are negatively correlated. Intuitively, a large number of tosses that result in a 1 suggests a smaller number of tosses that result in a 2.
- (b) Let  $A_t$  (respectively,  $B_t$ ) be a Bernoulli random variable that is equal to 1 if and only if the  $t$ th toss resulted in 1 (respectively, 2). We have  $\mathbf{E}[A_t B_t] = 0$  (since  $A_t \neq 0$  implies  $B_t = 0$ ) and

$$\mathbf{E}[A_t B_s] = \mathbf{E}[A_t] \mathbf{E}[B_s] = \frac{1}{k} \cdot \frac{1}{k} \quad \text{for } s \neq t.$$

Thus,

$$\begin{aligned} \mathbf{E}[X_1 X_2] &= \mathbf{E}[(A_1 + \dots + A_n)(B_1 + \dots + B_n)] \\ &= n \mathbf{E}[A_1(B_1 + \dots + B_n)] = n(n-1) \cdot \frac{1}{k} \cdot \frac{1}{k} \end{aligned}$$

and

$$\begin{aligned}
 \text{cov}(X_1, X_2) &= \mathbf{E}[X_1 X_2] - \mathbf{E}[X_1] \mathbf{E}[X_2] \\
 &= \frac{n(n-1)}{k^2} - \frac{n^2}{k^2} = -\frac{n}{k^2}.
 \end{aligned}$$

The covariance of  $X_1$  and  $X_2$  is negative as expected.

4. (a) If  $X$  takes a value  $x$  between  $-1$  and  $1$ , the conditional PDF of  $Y$  is uniform between  $-2$  and  $2$ . If  $X$  takes a value  $x$  between  $1$  and  $2$ , the conditional PDF of  $Y$  is uniform between  $-1$  and  $1$ .

Similarly, if  $Y$  takes a value  $y$  between  $-1$  and  $1$ , the conditional PDF of  $X$  is uniform between  $-1$  and  $2$ . If  $Y$  takes a value  $y$  between  $1$  and  $2$ , or between  $-2$  and  $-1$ , the conditional PDF of  $X$  is uniform between  $-1$  and  $1$ .

- (b) We have

$$\mathbf{E}[X | Y = y] = \begin{cases} 0, & \text{if } -2 \leq y \leq -1, \\ 1/2, & \text{if } -1 < y \leq 1, \\ 0, & \text{if } 1 \leq y \leq 2, \end{cases}$$

and

$$\text{var}(X | Y = y) = \begin{cases} 1/3, & \text{if } -2 \leq y \leq -1, \\ 3/4, & \text{if } -1 < y \leq 1, \\ 1/3, & \text{if } 1 \leq y \leq 2. \end{cases}$$

It follows that  $\mathbf{E}[X] = 3/10$  and  $\text{var}(X) = 193/300$ .

- (c) By symmetry, we have  $\mathbf{E}[Y | X] = 0$  and  $\mathbf{E}[Y] = 0$ . Furthermore,  $\text{var}(Y | X = x)$  is the variance of a uniform PDF (whose range depends on  $x$ ), and

$$\text{var}(Y | X = x) = \begin{cases} 4/3, & \text{if } -1 \leq x \leq 1, \\ 1/3, & \text{if } 1 < x \leq 2. \end{cases}$$

Using the law of total variance, we obtain

$$\text{var}(Y) = \mathbf{E}[\text{var}(Y | X)] = \frac{4}{5} \cdot \frac{4}{3} + \frac{1}{5} \cdot \frac{1}{3} = 17/15.$$

5. First let us write out the properties of all of our random variables. Let us also define  $K$  to be the number of members attending a meeting and  $B$  to be the Bernoulli random variable describing whether or not a member attends a meeting.

$$\begin{aligned}
 \mathbf{E}[N] &= \frac{1}{1-p}, & \text{var}(N) &= \frac{p}{(1-p)^2}, \\
 \mathbf{E}[M] &= \frac{1}{\lambda}, & \text{var}(M) &= \frac{1}{\lambda^2}, \\
 \mathbf{E}[B] &= q, & \text{var}(B) &= q(1-q).
 \end{aligned}$$

- (a) Since  $K = B_1 + B_2 + \cdots + B_N$ ,

$$\begin{aligned}
 \mathbf{E}[K] &= \mathbf{E}[N] \cdot \mathbf{E}[B] = \frac{q}{1-p}, \\
 \text{var}(K) &= \mathbf{E}[N] \cdot \text{var}(B) + (\mathbf{E}[B])^2 \cdot \text{var}(N) = \frac{q(1-q)}{1-p} + \frac{pq^2}{(1-p)^2}.
 \end{aligned}$$

(b) Let  $G$  be the total money brought to the meeting. Then  $G = M_1 + M_2 + \cdots + M_K$ .

$$\begin{aligned}
 \mathbf{E}[G] &= \mathbf{E}[M] \cdot \mathbf{E}[K] = \frac{q}{\lambda(1-p)}, \\
 \text{var}(G) &= \text{var}(M) \cdot \mathbf{E}[K] + (\mathbf{E}[M])^2 \text{var}(K) \\
 &= \frac{q}{\lambda^2(1-p)} + \frac{1}{\lambda^2} \left( \frac{q(1-q)}{1-p} + \frac{pq^2}{(1-p)^2} \right).
 \end{aligned}$$

G1<sup>†</sup>. (a) Let  $X_1, X_2, \dots, X_n$  be independent, identically distributed (IID) random variables. We note that

$$\mathbf{E}[X_1 + \cdots + X_n \mid X_1 + \cdots + X_n = x_0] = x_0.$$

It follows from the linearity of expectations that

$$\begin{aligned}
 x_0 &= \mathbf{E}[X_1 + \cdots + X_n \mid X_1 + \cdots + X_n = x_0] \\
 &= \mathbf{E}[X_1 \mid X_1 + \cdots + X_n = x_0] + \cdots + \mathbf{E}[X_n \mid X_1 + \cdots + X_n = x_0]
 \end{aligned}$$

Because the  $X_i$ 's are identically distributed, we have the following relationship.

$$\mathbf{E}[X_i \mid X_1 + \cdots + X_n = x_0] = \mathbf{E}[X_j \mid X_1 + \cdots + X_n = x_0], \text{ for any } 1 \leq i \leq n, 1 \leq j \leq n.$$

Therefore,

$$\begin{aligned}
 n\mathbf{E}[X_1 \mid X_1 + \cdots + X_n = x_0] &= x_0 \\
 \mathbf{E}[X_1 \mid X_1 + \cdots + X_n = x_0] &= \frac{x_0}{n}.
 \end{aligned}$$

(b) Note that we can rewrite  $\mathbf{E}[X_1 \mid S_n = s_n, S_{n+1} = s_{n+1}, \dots, S_{2n} = s_{2n}]$  as follows:

$$\begin{aligned}
 &\mathbf{E}[X_1 \mid S_n = s_n, S_{n+1} = s_{n+1}, \dots, S_{2n} = s_{2n}] \\
 &= \mathbf{E}[X_1 \mid S_n = s_n, X_{n+1} = s_{n+1} - s_n, X_{n+2} = s_{n+2} - s_{n+1}, \dots, X_{2n} = s_{2n} - s_{2n-1}] \\
 &= \mathbf{E}[X_1 \mid S_n = s_n],
 \end{aligned}$$

where the last equality holds due to the fact that the  $X_i$ 's are independent. We also note that

$$\mathbf{E}[X_1 + \cdots + X_n \mid S_n = s_n] = \mathbf{E}[S_n \mid S_n = s_n] = s_n.$$

It follows from the linearity of expectations that

$$\mathbf{E}[X_1 + \cdots + X_n \mid S_n = s_n] = \mathbf{E}[X_1 \mid S_n = s_n] + \cdots + \mathbf{E}[X_n \mid S_n = s_n].$$

Because the  $X_i$ 's are identically distributed, we have the following relationship:

$$\mathbf{E}[X_i \mid S_n = s_n] = \mathbf{E}[X_j \mid S_n = s_n], \text{ for any } 1 \leq i \leq n, 1 \leq j \leq n.$$

Therefore,

$$\mathbf{E}[X_1 + \cdots + X_n \mid S_n = s_n] = n\mathbf{E}[X_1 \mid S_n = s_n] = s_n \Rightarrow \mathbf{E}[X_1 \mid S_n = s_n] = \frac{s_n}{n}.$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.