

Problem Set 9 Solutions

Solution to Problem 1: Well, Well, Well

Solution to Problem 1, part a.

Inside the well $V(x) = 0$ and therefore

$$i\hbar \frac{\partial \psi(x, t)}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} \quad (9-6)$$

Solution to Problem 1, part b.

If E has a nonzero imaginary part E_{imag} , then the magnitude of $f(t)$ is a function of time, in particular

$$|f(t)| = \exp(E_{imag}t/\hbar) \quad (9-7)$$

If $E_{imag} > 0$ then $|f(t)|$ gets large for large values of t (i.e., it blows up at infinity). If $E_{imag} < 0$ then $|f(t)|$ gets large for large values of $-t$ (i.e., it blows up at negative infinity). In either case it is impossible to normalize $\psi(x)$.

Solution to Problem 1, part c.

$$E\phi(x) = -\frac{\hbar^2}{2m} \frac{\partial^2 \phi(x, t)}{\partial x^2} \quad (9-8)$$

Solution to Problem 1, part d.

Since

$$\phi(x) = a \sin(kx) + b \cos(kx) \quad (9-9)$$

$$\frac{d\phi(x)}{dx} = ak \cos(kx) - bk \sin(kx) \quad (9-10)$$

$$\begin{aligned} \frac{d^2\phi(x)}{dx^2} &= -ak^2 \sin(kx) - bk^2 \cos(kx) \\ &= -k^2 \phi(x) \end{aligned} \quad (9-11)$$

Therefore

$$E\phi(x) = \left(\frac{\hbar^2 k^2}{2m} \right) \phi(x) \quad (9-12)$$

so

$$E = \frac{\hbar^2 k^2}{2m} \quad (9-13)$$

Solution to Problem 1, part e.

One of the boundary conditions is $\phi(0) = 0$, so

$$\begin{aligned} 0 &= \phi(0) \\ &= a\sin(0) + b\cos(0) \\ &= b \end{aligned} \tag{9-14}$$

Since we know the wavefunction is nonzero, a must be nonzero as well.

Solution to Problem 1, part f.

$\phi(x)$ must be zero at the boundaries, which implies

$$\frac{k = j\pi}{L} \tag{9-15}$$

so that $\sin(-kL) = 0$.

Solution to Problem 1, part g.

$$e_j = \frac{\hbar^2 \pi^2 j^2}{2mL^2} \tag{9-16}$$

Solution to Problem 1, part h.

$$\phi_j(x) = a \sin\left(\frac{j\pi x}{L}\right) \tag{9-17}$$

Solution to Problem 1, part i.

$$e_1 = \frac{\hbar^2 \pi^2}{2mL^2} \tag{9-18}$$

Solution to Problem 1, part j.

$$e_2 = \frac{2\hbar^2 \pi^2}{mL^2} \tag{9-19}$$

Solution to Problem 1, part k.

$$e_1 = \frac{\hbar^2 \pi^2}{2mL^2} \tag{9-20}$$

$$= \frac{(1.054 \times 10^{-34} \text{Joule-seconds})^2 \times (3.1416)^2}{2 \times (9.109 \times 10^{-31} \text{kilograms}) \times (2 \times 10^{-8} \text{meters})^2} \tag{9-21}$$

$$= 1.506 \times 10^{-22} \text{Joules} \tag{9-22}$$

$$\tag{9-23}$$

Solution to Problem 1, part l.

Express this ground-state energy in electron-volts (1 eV = 1.602×10^{-19} Joules).

$$\begin{aligned} e_1 &= 1.506 \times 10^{-22} \text{Joules} \\ &= 9.391 \times 10^{-4} \text{eV} \end{aligned} \tag{9-24}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

6.050J / 2.110J Information and Entropy
Spring 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.