

$$\frac{d}{dx}a^x?$$

We now want to learn to differentiate *any* exponential  $a^x$ . There are two roughly equivalent methods we can use:

**Method 1: Convert  $a^x$  to something with base e and use the chain rule.**

Because  $\ln x$  is the inverse function to  $e^x$  we can rewrite  $a$  as  $e^{\ln(a)}$ . Thus:

$$a^x = \left(e^{\ln(a)}\right)^x = e^{x \ln(a)}$$

That looks like it might be tricky to differentiate. Let's work up to it:

$$\begin{aligned}\frac{d}{dx}e^x &= e^x \\ \frac{d}{dx}e^{3x} &= 3e^{3x} \quad (\text{by the chain rule})\end{aligned}$$

Remember,  $\ln(a)$  is just a constant like 3, not a variable. Therefore:

$$\frac{d}{dx}e^{(\ln a)x} = (\ln a)e^{(\ln a)x}$$

or

$$\frac{d}{dx}a^x = (\ln a)a^x$$

This is a common type of calculation; you should practice it until you are comfortable with it. You may either memorize formulas for  $\frac{d}{dx}e^{kx}$  and  $\frac{d}{dx}a^x$  or re-derive them every time you need them.

Recall that  $\frac{d}{dx}a^x = M(a) \cdot a^x$ . So finally we know the value of  $M(a)$ :

$$M(a) = \ln(a)$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.01SC Single Variable Calculus  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.