

8.286 Lecture 11
October 17, 2013

NON-EUCLIDEAN SPACES: CLOSED UNIVERSES

REVIEW OF LECTURE 10: INTRODUCTION TO NON-EUCLIDEAN SPACES


-1-

curved space?

curves with respect to what?

curved spacetime?

non-Euclidean geometry



Note on image: I have added the "Dali" clock and Fermi satellite images to the original image created for the Gravity Probe B collaboration
Slides courtesy of Mustafa Amin. Used with permission.

-2-

Euclid's *Postulates*




1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given a straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
4. All right angles are congruent.
5. If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines if produced indefinitely meet on that side on which the angles are less than two right angles

Corrected 10/10/13

Slides courtesy of Mustafa Amin. Used with permission.

-3-

5th Postulate




5. If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines if produced indefinitely meet on that side on which the angles are less than two right angles

Corrected 10/10/13


Slides courtesy of Mustafa Amin. Used with permission.

Equivalent Statements of the 5th Postulate




(a) "If a straight line intersects one of two parallels (i.e., lines which do not intersect however far they are extended), it will intersect the other also."

Equivalent Statements of the 5th Postulate




(b) "There is one and only one line that passes through any given point and is parallel to a given line."

Equivalent Statements of the 5th Postulate




(c) "Given any figure there exists a figure, similar to it, of any size." (Two polygons are similar if their corresponding angles are equal, and their corresponding sides are proportional.)

Equivalent Statements of the 5th Postulate



(d) "There is a triangle in which the sum of the three angles is equal to two right angles (i.e., 180°)."

Giovanni Gerolamo Saccheri (1667-1733)



In 1733, Saccheri, a Jesuit priest, published *Euclides ab omni naevo vindicatus* (*Euclid Freed of Every Flaw*).

The book was a study of what geometry would be like if the 5th postulate were false.

He hoped to find an inconsistency, but failed.

© Source unknown. All rights reserved. This content is excluded from our Creative Commons license. For more information, see <http://ocw.mit.edu/fairuse>.

Carl Friedrich Gauss (1777-1855)



German mathematician and physicist.

Born as the son of a poor working-class parents. His mother was illiterate and never even recorded the date of his birth.

His students included Richard Dedekind, Bernhard Riemann, Peter Gustav Lejeune Dirichlet, Gustav Kirchhoff, and August Ferdinand Möbius.

Photo from http://commons.wikimedia.org/wiki/File:Carl_Friedrich_Gauss.jpg (in public domain).


~1750-1850



- infinite
- constant negative curvature



~~5th Postulate~~



Slides courtesy of Mustafa Amin. Used with permission.

GBL geometry with Klein



1. constant negative curvature
2. infinite
3. ~~5th postulate~~

(x_1, y_1)

$$x^2 + y^2 < 1$$


(x_2, y_2)

$$d(1,2) = a \cosh^{-1} \left[\frac{1 - x_1 x_2 - y_1 y_2}{\sqrt{1 - x_1^2 - y_1^2} \sqrt{1 - x_2^2 - y_2^2}} \right]$$

Note: no global embedding in 3D Euclidean space possible

-12-

Geometry (after Klein)



co-ordinates

(x_1, y_1)



(x_2, y_2)

distance function

$$d[(x_1, y_1), (x_2, y_2)]$$

-13-

tiny distances




$$ds^2 = dx^2 + dy^2$$

$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

-14-


quadratic form



$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

-15-

locally Euclidean



$$ds^2 = dx^2 + dy^2$$


$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

$$g_{xx}g_{yy} - g_{xy}^2 > 0$$

Slides courtesy of Mustafa Amin. Used with permission.

-16-


Non-Euclidean Geometry: The Surface of a Sphere



$$x^2 + y^2 + z^2 = R^2 .$$


-17-

Polar Coordinates:




-18-

Varying θ :




-19-

Varying ϕ :



A Sphere in 4 Euclidean Dimensions



MIT OpenCourseWare
<http://ocw.mit.edu>

8.286 The Early Universe
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.